WARNINGS:

- · Handle with care.
- · Skin and eye irritant.
- · Store in a cool place, away from food, feeds and seed.
- · Keep out of reach of children, uninformed persons and animals.
- Full cognizance must be taken to all WARNINGS, PRECAUTIONS and DIRECTIONS FOR USE mentioned on the label of the products with which TYBOOST will be used.

PRECAUTIONS:

- Do not inhale fumes or spray mist.
- Wash with soap and water after use and accidental skin contact.
- Do not eat, drink or smoke while mixing or spraving unless having washed hands and face.
- TYBOOST may be corrosive to metal parts of sprayers.
- Thoroughly wash and rinse spray equipment after use and dispose of wash water where it will not contaminate crops, grazing, rivers and dams.
- TRIPLE RINSE empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the contents of the spray tank before destroying the container in the prescribed manner.
- Do not re-use the empty container for any other purpose but destroy it by perforation and flattening and dispose of it in a safe way.
- Prevent contamination of food, feedstuffs. drinking water and eating utensils.

TYBOOST

Reg. No. L 10834 Act/Wet No. 36 of/van 1947

An aqueous solution salt adjuvant to improve the uptake of the herbicides listed

'n Wateroplosbare sout bevorderingsmiddel om die opneembaarheid van die genoemde onkruiddoders te verhoog.

ACTIVE INGREDIENT: / AKTIEWE BESTANDDEEL:

ammonium sulphate	500	g/l
ammoniumsulfaat	500	g/l

Registration holder / Registrasiehouer: GREEN ISLAND INVESTMENTS (Pty) Ltd.

Co. Reg. No./Mpy. Reg. Nr.: 2013/043963/07 P.O. Box / Posbus 1822, Heidelberg 1438, South Africa. Tel: +27 10 035 4143

ADJUVANT

CAUTION

VERSIGTIG

Batch No./Lot Nr.: Date formulated: Formuleringsdatum: U.N. No. 3082





Net Volume:

Netto Volume:



p**hoo**n

DIRECTIONS FOR USE: USE ONLY AS DIRECTED. SHAKE WELL BEFORE USE

General Information:

- **TYBOOST** improves the performance of herbicides which are sensitive to salt antagonism such as glycines, phenoxy-carboxylic acids (eg 2.4-D. MCPA), benzoic acids, cvclohexanediones, aryloxyphenoxy propionates and imidazolinone herbicides.
- Always refer to the relevant herbicide label regarding sensitivity to salt antagonism, acidification and instructions for use.
- Tank mix **TYBOOST** with the relevant herbicide and apply as a post-emergent application to weeds.
- Use TYBOOST at a rate of 2% (volume/volume) of the sprav mixture.
- Ensure thorough coverage and wetting of the weeds, especially when crop foliage is dense.

Mixing Instructions:

READ THE LABEL IN DETAIL BEFORE OPENING THE CONTAINER / LEES DIE ETIKET VOLLEDIG VOORDAT DIE HOUEN OOPGEMAAK WORD.

- Three quarter fill the spray tank with clean water.
- Measure the required volume of **TYBOOST** and pre-mix this with at least 10^ℓ of water before adding to the spray tank.
- The other product(s) to be mixed with TYBOOST should be pre-mixed in similar fashion.
- Agitate the water in the spray tank and then first add the **TYBOOST** followed by the other product(s).
- Fill the sprav tank with water to the required level while maintaining agitation to ensure thorough mixing of the spray mixture before spraying commences. Maintain agitation while spraying.
- The prepared spray mixture must not be left in the spray tank for any length of time, e.g. overnight.

All spray applications must be made with suitable equipment that is in good working order and correctly calibrated to give the desired coverage for that particular method of application.

DOSAGE RATE FOR APPLICATION IN TANK MIXTURES WITH SPECIFIED HERBICIDE: Mix 2 ℓ **TYBOOST** per 100 ℓ water (2%).